

5

Local Phase Invariance and Electrodynamics

The theories that make up the standard model are all based on the principle of local symmetry. The simplest example of a local symmetry is the extension of the global phase invariance discussed at the end of Note 2 to local phase invariance. As we will derive below, the requirement that a theory be invariant under local phase transformations implies the existence of a gauge field in the theory that mediates or carries the "force" between the matter fields. For electrodynamics the gauge field is the electromagnetic vector potential $A_\mu(x)$ and its quantum particle is the massless photon. In addition, in the standard model the gauge fields mediating the strong interactions between the quarks are the massless gluon fields and the gauge fields mediating the weak interactions are the fields for the massive Z^0 and W^\pm weak bosons.

To illustrate these principles we extend the global phase invariance of the Lagrangian of Eq. 1 to a theory that has local phase invariance. Thus, we require \mathcal{L} to have the same form for $\varphi'(x)$ and $\varphi(x)$, where the local phase transformation is defined by

$$\varphi'(x) = e^{ie(x)}\varphi(x). \quad (30)$$

The potential energy,

$$V(\varphi, \varphi^\dagger) = m^2\varphi^\dagger\varphi + \lambda(\varphi^\dagger\varphi)^2, \quad (31)$$

already has this symmetry, but the kinetic energy, $\partial^\mu\varphi^\dagger\partial_\mu\varphi$, clearly

does not, since

$$\partial_\mu\varphi'(x) = e^{ie(x)}[\partial_\mu\varphi + i(\partial_\mu\epsilon)\varphi]. \quad (32)$$

\mathcal{L} does not have local phase invariance if the Lagrangian of the transformed fields depends on $\epsilon(x)$ or its derivatives. The way to eliminate the $\partial_\mu\epsilon$ dependence is to add a new field $A_\mu(x)$ called the gauge field and then require the local symmetry transformation law for this new field to cancel the $\partial_\mu\epsilon$ term in Eq. 32. The gauge field can be added by generalizing the derivative ∂_μ to D_μ , where

$$D_\mu = \partial_\mu - ieA_\mu(x). \quad (33)$$

This is just the minimal-coupling procedure of electrodynamics. We can then make a kinetic energy term of the form $(D^\mu\varphi)^\dagger(D_\mu\varphi)$ if we require that

$$D'_\mu\varphi'(x) = e^{ie(x)}D_\mu\varphi(x). \quad (34)$$

When written out with Eq. 33, Eq. 34 becomes an equation for $A'_\mu(x)$ in terms of $A_\mu(x)$, which is easily solved to give

$$A'_\mu(x) = A_\mu(x) + \frac{1}{e}\partial_\mu\epsilon(x). \quad (35)$$

Equation 35 prescribes how the gauge field transforms under the local phase symmetry.

Thus the first step to modifying Eq. 1 to be a theory with local phase invariance is simply to replace ∂_μ by D_μ in \mathcal{L} . (A slightly generalized form of this trick is used in the construction of all the theories in the standard model.) With this procedure the dominant interaction of the gauge field $A^\mu(x)$ with the matter field φ is in the form of a current times the gauge field, $eJ^\mu A_\mu$, where J_μ is the current defined in Eq. 14.

6

Spontaneous Breaking of Local Phase Invariance

We now show that spontaneous breaking of local symmetry implies that the associated vector boson has a mass, in spite of the fact that $A^\mu A_\mu$ by itself is not locally phase invariant. Much of the calculation in Note 3 can be translated to the Lagrangian of Eq. 38. In fact, the calculation is identical from Eq. 16 to Eq. 18, so the first new step is to substitute Eq. 17 into Eq. 38. The only significantly new part

of the calculation is replacing $\partial^\mu\varphi^\dagger\partial_\mu\varphi$ by $(D^\mu\varphi)^\dagger(D_\mu\varphi)$. However, instead of simply substituting Eq. 17 for φ and computing $(D^\mu\varphi)^\dagger(D_\mu\varphi)$ directly, it is convenient to make a local phase transformation first:

$$\varphi'(x) = \frac{1}{\sqrt{2}}[\rho(x) + \varphi_0] \exp[i\pi(x)/\varphi_0], \quad (41)$$

where $\varphi(x) = [\rho(x) + \varphi_0]/\sqrt{2}$. (The local phase invariance permits us to remove the phase of $\varphi(x)$ at every space-time point.) We emphasize the difference between Eqs. 17 and 41: Eq. 17 defines the $\rho(x)$ and $\pi(x)$ fields; Eq. 41 is a local phase transformation of $\varphi(x)$ by angle $\pi(x)$. Don't be fooled by the formal similarity of the two equations. Thus, we may write Eq. 38 in terms of $\varphi(x) = [\rho(x) + \varphi_0]/\sqrt{2}$ and obtain

This leaves a problem. If we simply replace $\partial_\mu\phi$ by $D_\mu\phi$ in the Lagrangian and then derive the equations of motion for A_μ , we find that A_μ is proportional to the current J_μ . The A_μ field equation has no space-time derivatives and therefore $A_\mu(x)$ does not propagate. If we want A_μ to correspond to the electromagnetic field potential, we must add a kinetic energy term for it to \mathcal{L} .

The problem then is to find a locally phase invariant kinetic energy term for $A_\mu(x)$. Note that the combination of covariant derivatives $D_\mu D_\nu - D_\nu D_\mu$, when acting on any function, contains no derivatives of the function. We define the electromagnetic field tensor of electrodynamics as

$$F_{\mu\nu} \equiv \frac{i}{e} [D_\mu, D_\nu] = \partial_\mu A_\nu - \partial_\nu A_\mu. \quad (36)$$

It contains derivatives of A_μ . Its transformation law under the local symmetry is

$$F'_{\mu\nu} = F_{\mu\nu}. \quad (37)$$

Thus, it is completely trivial to write down a term that is quadratic in the derivatives of A_μ , which would be an appropriate kinetic energy term. A fully phase invariant generalization of Eq. 1a is

$$\mathcal{L} = -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} + (D^\mu\phi)^\dagger (D_\mu\phi) - m^2\phi^\dagger\phi - \lambda(\phi^\dagger\phi)^2. \quad (38)$$

We should emphasize that \mathcal{L} has no mass term for $A_\mu(x)$. Thus, when the fields correspond directly to the particles in Eq. 38, the vector particles described by $A_\mu(x)$ are massless. In fact, $A^\mu A_\mu$ is not invariant under the gauge transformation in Eq. 35, so it is not obvious how the A_μ field can acquire a mass if the theory does have local phase invariance. In Note 6 we will show how the gauge field becomes massive through spontaneous symmetry breaking. This is

the key to understanding the electroweak theory.

We now rediscover the Lagrangian of electrodynamics for the interaction of electrons and photons following the same procedure that we used for the complex scalar field. We begin with the kinetic energy term for a Dirac field of the electron ψ , replace ∂_μ by D_μ defined in Eq. 33, and then add $-\frac{1}{4}F^{\mu\nu}F_{\mu\nu}$, where $F^{\mu\nu}$ is defined in Eq. 36. The Lagrangian for a free Dirac field is

$$\mathcal{L}_{\text{Dirac}} = \bar{\psi}(i\gamma^\mu\partial_\mu - m)\psi, \quad (39)$$

where γ^μ are the four Dirac γ matrices and $\bar{\psi} = \psi^\dagger\gamma_0$. Straightening out the definition of the γ^μ matrices and the components of ψ is the problem of describing a spin-1/2 particle in a theory with Lorentz invariance. We leave the details of the Dirac theory to textbooks, but note that we will use some of these details when we finally write down the interactions of the quarks and leptons. The interaction of the electron field ψ with the electromagnetic field follows by replacing ∂_μ by D_μ . The electrodynamic Lagrangian is

$$\mathcal{L} = -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} + \bar{\psi}(i\gamma^\mu D_\mu - m)\psi, \quad (40a)$$

where the interaction term in $i\bar{\psi}\gamma^\mu D_\mu\psi$ has the form

$$\mathcal{L}_{\text{interaction}} = e\bar{\psi}\gamma_\mu\psi A^\mu = eJ_\mu^{\text{em}} A^\mu, \quad (40b)$$

where $J_\mu^{\text{em}} = \bar{\psi}\gamma_\mu\psi$ is the electromagnetic current of the electron. What is amazing about the standard model is that all the electroweak and strong interactions between fermions and vector bosons are similar in form to Eq. 40b, and much phenomenology can be understood in terms of such interaction terms as long as we can approximate the quantum fields with the classical solutions.

$$\mathcal{L} = -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} + \frac{1}{2} \partial^\mu\rho\partial_\mu\rho + \frac{e^2}{2}(\rho + \phi_0)^2 A^\mu A_\mu - \frac{m^2}{2}(\rho + \phi_0)^2 - \frac{\lambda}{4}(\rho + \phi_0)^4. \quad (42)$$

(At the expense of a little algebra, the calculation can be done the other way. First substitute Eq. 17 for ϕ in Eq. 38. One then finds an $A^\mu\partial_\mu\pi$ term in \mathcal{L} that can be removed using the local phase transformation $A'_\mu = A_\mu - [1/(e\phi_0)]\partial_\mu\pi$, $\rho' = \rho$, and $\pi' = 0$. Equation 42 then follows, although this method requires some effort. Thus, a reason for doing the calculation in the order of Eq. 41 is that the algebra gets messy rather quickly if the local symmetry is not used early in the calculation of the electroweak case. However, in principle it makes little difference.)

The Lagrangian in Eq. 42 is an amazing result: the π field has

vanished from \mathcal{L} altogether (according to Eq. 41, it was simply a gauge artifact), and there is a term $\frac{1}{2}e^2\phi_0^2 A^\mu A_\mu$ in \mathcal{L} , which is a mass term for the vector particle. Thus, the massless particle of the global case has become the longitudinal mode of a massive vector particle, and there is only one scalar particle ρ left in the theory. In somewhat more picturesque language the vector boson has eaten the Goldstone boson and become heavy from the feast. However, the existence of the vector boson mass terms should not be understood in isolation: the phase invariance of Eq. 42 determines the form of the interaction of the massive A_μ field with the ρ field.

This calculation makes it clear that it can be tricky to derive the spectrum of a theory with local symmetry and spontaneous symmetry breaking. Theoretical physicists have taken great care to confirm that this interpretation is correct and that it generalizes to the full quantum field theory.

7

Yang-Mills Theories

The standard model possesses symmetries of the type described in Note 4, except that they are local. Thus, we need to carry out the calculations of Note 5 for Lie-group symmetries. As the reader might expect, this requires replacing $\epsilon(x)$ of Eq. 13 by a matrix or, equivalently, the matrix of Eq. 21 by a matrix function of x , $\epsilon^a(x)T_a$. The Yang-Mills Lagrangian can be derived by mimicking with matrix functions Eqs. 34 to 38.

The internal, local transformation of the ϕ field (ϕ is a column vector with components ϕ_i , where i runs from 1 to n) is

$$\phi'(x) = e^{i\epsilon(x)}\phi(x), \quad (43)$$

which is formally identical to Eq. 30, except that $\epsilon(x)$ is now an n -by- n matrix. Thus,

$$\epsilon(x) = \epsilon^a(x)T_a, \quad (44)$$

where the sum on a is over the N independent symmetries. Equation 43 is a symmetry of the potential energy

$$V = \mu^2\phi^\dagger\phi + \lambda(\phi^\dagger\phi)^2, \quad (45)$$

if $\epsilon(x)$ in Eq. 44 is a Hermitian matrix (that is, if $T_a = T_a^\dagger$ and the $\epsilon^a(x)$ are real functions). The kinetic energy $(\partial^\mu\phi)^\dagger(\partial_\mu\phi)$ can be made phase invariant by extending ∂_μ to D_μ , analogous to Eq. 33 for electrodynamics:

$$D_\mu = \partial_\mu - ieA_\mu, \quad (46a)$$

where

$$A_\mu = A_\mu^a T_a, \quad (46b)$$

so that A_μ is an n -by- n matrix that acts on the ϕ vector. Just as for Eq. 35, the transformation properties of A_μ are derived from the equation

$$D'_\mu\phi'(x) = e^{i\epsilon(x)}D_\mu\phi(x). \quad (47)$$

After some matrix manipulation one finds the solution of Eq. 47 for $A'_\mu(x)$ in terms of $A_\mu(x)$ to be

$$A'_\mu(x) = e^{i\epsilon(x)}A_\mu(x)e^{-i\epsilon(x)} - \frac{1}{e}\partial_\mu\epsilon(x), \quad (48)$$

where $e^{-i\epsilon(x)}$ is the inverse of the matrix $e^{i\epsilon(x)}$. With these requirements, it is easily seen that $(D^\mu\phi)^\dagger(D_\mu\phi)$ is invariant under the group of local transformations.

The calculation of the field tensor is formally identical to Eq. 36, except we must take into account that $A_\mu(x)$ is a matrix. Thus, we define a matrix $F_{\mu\nu}$ field tensor as

$$F_{\mu\nu} \equiv \frac{i}{e}[D_\mu, D_\nu] = \partial_\mu A_\nu - \partial_\nu A_\mu - ie[A_\mu, A_\nu]. \quad (49)$$

There is a field tensor for each group generator, and some further matrix manipulation plus Eq. 26 gives the components,

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + ef^{abc}A_{\mu b}A_{\nu c}. \quad (50)$$

The transformation law for the matrix $F_{\mu\nu}$ is

$$F'_{\mu\nu} = e^{i\epsilon(x)}F_{\mu\nu}e^{-i\epsilon(x)}. \quad (51)$$

Thus, we can write down a kinetic energy term in analogy to electrodynamics:

$$\mathcal{L}_{\text{kinetic energy}} = -\frac{1}{4}F_{\mu\nu}^a F_a^{\mu\nu}. \quad (52)$$

The locally invariant Yang-Mills Lagrangian for spinless fields coupled to the vector bosons is


$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^a F_a^{\mu\nu} + (D^\mu\phi)^\dagger(D_\mu\phi) - \mu^2\phi^\dagger\phi - \lambda(\phi^\dagger\phi)^2. \quad (53)$$

Just as in electrodynamics, we can add fermions to the theory in the form

$$\mathcal{L}_{\text{fermion}} = \bar{\psi}(i\gamma^\mu D_\mu - m)\psi, \quad (54)$$

where D_μ is defined in Eq. 46 and ψ is a column vector with n_f entries ($n_f =$ number of fermions). The matrices T_a in D_μ for the fermion covariant derivative are usually different from the matrices for the spinless fields, since there is no requirement that ϕ and ψ need to belong to the same representation of the group. It is, of course, necessary for the sets of T_a matrices to satisfy the commutation relations of Eq. 26 with the same set of structure constants.

We will not look at the general case of spontaneous symmetry breaking in a Yang-Mills theory, which is a messy problem mathematically. There is spontaneous symmetry breaking in the electroweak sector of the standard model, and we will work out the steps analogous to Eqs. 41 and 42 for this particular case in the next Note.



The $SU(2) \times U(1)$ Electroweak Model

The main emphasis in these Notes has been on developing just those aspects of Lagrangian field theory that are needed for the standard model. We have now come to the crucial step: finding a Lagrangian that describes the electroweak interactions. It is rather difficult to be systematic. The historical approach would be complicated by the rather late discovery of the weak neutral currents, and a purely phenomenological development is not yet totally logical because there are important aspects of the standard model that have not yet been tested experimentally. (The most important of these are the details of the spontaneous symmetry breaking.) Although we will write down the answer without excessive explanation, the reader should not forget the critical role that experimental data played in the development of the theory.

The first problem is to identify the local symmetry group. Before the standard model was proposed over twenty years ago, the electromagnetic and charge-changing weak interactions were known. The smallest continuous group that can describe these is $SU(2)$, which has a doublet representation. If the weak interactions can change electrons to electron neutrinos, which are electrically neutral, it is not possible to incorporate electrodynamics in $SU(2)$ alone unless a heavy positively charged electron is added to the electron and its neutrino to make a triplet, because the sum of charges in an $SU(2)$ multiplet is zero. Various schemes of this sort have been tried but do not agree with experiment. The only way to leave the electron and electron neutrino in a doublet and include electrodynamics is to add an extra $U(1)$ interaction to the theory. The hypothesis of the extra $U(1)$ factor was challenged many times until the discovery of the weak neutral current. That discovery established that the local symmetry of the electroweak theory had to be at least as large as $SU(2) \times U(1)$.

Let us now interpret the physical meaning of the four generators of $SU(2) \times U(1)$. The three generators of the $SU(2)$ group are I^+ , I_3 , and I^- , and the generator of the $U(1)$ group is called Y , the weak hypercharge. (The weak $SU(2)$ and $U(1)$ groups are distinguished from other $SU(2)$ and $U(1)$ groups by the label “W.”) I^+ and I^- are associated with the weak charge-changing currents (the general definition of a current is described in Note 2), and the charge-changing currents couple to the W^+ and W^- charged weak vector bosons in analogy to Eq. 40b. Both I_3 and Y are related to the electromagnetic current and the weak neutral current. In order to assign the electron and its neutrino to an $SU(2)$ doublet, the electric charge Q^{em} is defined by

$$Q^{em} = I_3 + Y/2, \tag{55}$$

so the sum of electric charges in an n -dimensional multiplet is $nY/2$. The charge of the weak neutral current is a different combination of I_3 and Y , as will be described below.

The Lagrangian includes many pieces. The kinetic energies of the vector bosons are described by \mathcal{L}_{Y-M} , in analogy to the first term in Eq. 38. The three weak bosons have masses acquired through spontaneous symmetry breaking, so we need to add a scalar piece $\mathcal{L}_{\text{scalar}}$ to the Lagrangian in order to describe the observed symmetry breaking (also see Eq. 38). The fermion kinetic energy $\mathcal{L}_{\text{fermion}}$ includes the fermion-boson interactions, analogous to the electromagnetic interactions derived in Eqs. 39 and 40. Finally, we can add terms that couple the scalars with the fermions in a term $\mathcal{L}_{\text{Yukawa}}$. One physical significance of the Yukawa terms is that they provide for masses of the quarks and charged leptons.

The standard model is then a theory with a very long Lagrangian with many fields. The electroweak Lagrangian has the terms

$$\mathcal{L}_{\text{electroweak}} = \mathcal{L}_{Y-M} + \mathcal{L}_{\text{scalar}} + \mathcal{L}_{\text{fermion}} + \mathcal{L}_{\text{Yukawa}}. \tag{56}$$

(The reader may find this construction to be ad hoc and ugly. If so, the motivation will be clear for searching for a more unified theory from which this Lagrangian can be derived. However, it is important to remember that, at present, the standard model is the pinnacle of success in theoretical physics and describes a broader range of natural phenomena than any theory ever has.)

The Yang-Mills kinetic energy term has the form given by Eq. 52 for the $SU(2)$ bosons, plus a term for the $U(1)$ field tensor similar to electrodynamics (Eqs. 36 and 38).

$$\mathcal{L}_{Y-M} = -\frac{1}{4} F_{\mu\nu}^a F_{\mu\nu}^a - \frac{1}{4} F^{\mu\nu} F_{\mu\nu}, \tag{57}$$

where the $U(1)$ field tensor is

$$F_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu \tag{58}$$

and the $SU(2)$ Yang-Mills field tensor is

$$F_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g\epsilon_{abc} W_\mu^b W_\nu^c, \tag{59}$$

where the ϵ_{abc} are the structure constants for $SU(2)$ defined in Eq. 24 and the W_μ^a are the Yang-Mills fields.

8 continued

$SU(2) \times U(1)$ has two factors, and there is an independent coupling constant for each factor. The coupling for the $SU(2)$ factor is called g , and it has become conventional to call the $U(1)$ coupling $g'/2$. The two couplings can be written in several ways. The $U(1)$ of electrodynamics is generated by a linear combination of I_3 and Y , and the coupling is, as usual, denoted by e . The other coupling can then be parameterized by an angle θ_w . The relations among g , g' , e , and θ_w are

$$e \equiv gg'/\sqrt{g^2+g'^2} \text{ and } \tan \theta_w \equiv g'/g. \quad (60)$$

These definitions will be motivated shortly. In the electroweak theory both couplings must be evaluated experimentally and cannot be calculated in the standard model.

The scalar Lagrangian requires a choice of representation for the scalar fields. The choice requires that the field with a nonzero vacuum value is electrically neutral, so the photon remains massless, but it must carry nonzero values of I_3 and Y so that the weak neutral boson (the Z_μ^0) acquires a mass from spontaneous symmetry breaking. The simplest assignment is

$$\varphi = \begin{pmatrix} \varphi^+ \\ \varphi^0 \end{pmatrix} \text{ and } \varphi^\dagger = (-\varphi^-, (\varphi^0)^\dagger), \quad (61)$$

where φ^+ has $I_3 = 1/2$ and $Y = 1$, and φ^0 has $I_3 = -1/2$ and $Y = 1$. Since φ does not have $Y = -1$ fields, it is necessary to make φ a complex doublet, so $(\varphi^+)^\dagger = -\varphi^-$ has $I_3 = -1/2$ and $Y = -1$, and $(\varphi^0)^\dagger$ has $I_3 = 1/2$ and $Y = -1$. Then we can write down the Lagrangian of the scalar fields as

$$\mathcal{L}_{\text{scalar}} = (D^\mu \varphi)^\dagger (D_\mu \varphi) - m^2 \varphi^\dagger \varphi - \lambda (\varphi^\dagger \varphi)^2, \quad (62)$$

where

$$D_\mu \varphi = \partial_\mu \varphi - i \frac{g'}{2} B_\mu \varphi - i \frac{g}{2} \tau_a W_\mu^a \varphi \quad (63)$$

is the covariant derivative. The 2-by-2 matrices τ_a are the Pauli matrices. The factor of $1/2$ is required because the doublet representation of the $SU(2)$ generators is $\tau_a/2$. The factor of $1/2$ in the B_μ term is due to the convention that the $U(1)$ coupling is $g'/2$ and the

assignment that the φ doublet has $Y = 1$. After the spontaneous symmetry breaking, three of the four scalar degrees of freedom are "eaten" by the weak bosons. Thus just one scalar escapes the feast and should be observable as an independent neutral particle, called the Higgs particle. It has not (?) yet been observed experimentally, and it is perhaps the most important particle in the standard model that does not yet have a firm phenomenological basis. (The minimum number of scalar fields in the standard model is four. Experimental data could eventually require more.)

We now carry out the calculation for the spontaneous symmetry breaking of $SU(2) \times U(1)$ down to the $U(1)$ of electrodynamics. Just as in the example worked out in Note 6, spontaneous symmetry breaking occurs when $m^2 < 0$ in Eq. 62. In contrast to the simpler case, it is rather important to set up the problem in a clever way to avoid an inordinate amount of computation. As in Eq. 41, we write the four degrees of freedom in the complex scalar doublet so that it looks like a local symmetry transformation times a simple form of the field:

$$\varphi(x) = \exp[i\pi^a(x)\tau_a/2\varphi_0] \begin{pmatrix} 0 \\ [\rho(x) + \varphi_0]/\sqrt{2} \end{pmatrix}. \quad (64)$$

We can then write the scalar fields in a new gauge where the phases of $\varphi(x)$ are removed:

$$\varphi'(x) = \exp[-i\pi^a(x)\tau_a/2\varphi_0]\varphi(x) = \begin{pmatrix} 0 \\ [\rho(x) + \varphi_0]/\sqrt{2} \end{pmatrix}, \quad (65)$$

where we have used the freedom of making local symmetry transformations to write $\varphi'(x)$ in a very simple form. This choice, called the unitary gauge, will make it easy to write out Eq. 63 in explicit matrix form. Let us drop all primes on the fields in the unitary gauge and redefine W_μ^a by the equation

$$\tau_a W_\mu^a = \begin{pmatrix} W_\mu^3 & W_\mu^1 - iW_\mu^2 \\ W_\mu^1 + iW_\mu^2 & -W_\mu^3 \end{pmatrix} = \begin{pmatrix} W_\mu^3 & \sqrt{2} W_\mu^+ \\ \sqrt{2} W_\mu^- & -W_\mu^3 \end{pmatrix}, \quad (66)$$

where the definition of the Pauli matrices is used in the first step, and the W^\pm fields are defined in the second step with a numerical factor that guarantees the correct normalization of the kinetic energy of the charged weak vector bosons.

Next, we write out the $D_\mu \varphi$ in explicit matrix form, using Eqs. 63, 65, and 66:

$$D_\mu \varphi = \frac{1}{\sqrt{2}} \begin{pmatrix} -i\sqrt{2}gW_\mu^+(\rho + \varphi_0)/2 \\ \partial_\mu \rho - i(g'B_\mu - gW_\mu^3)(\rho + \varphi_0)/2 \end{pmatrix}. \quad (67)$$

Finally, we substitute Eqs. 65 and 67 into Eq. 63 and obtain

$$\begin{aligned} \mathcal{L}_{\text{scalar}} = & \frac{g^2}{4} W^\pm W_\mu^\pm (\rho + \varphi_0)^2 + \frac{1}{2} \partial^\mu \rho \partial_\mu \rho \\ & + \frac{1}{8} (g' B^\mu - g W_\mu^3) (g' B_\mu - g W_\mu^3) (\rho + \varphi_0)^2 \\ & + \frac{m^2}{2} (\rho + \varphi_0)^2 + \frac{\lambda}{4} (\rho + \varphi_0)^4, \end{aligned} \quad (68)$$

where ρ is the, as yet (?), unobserved Higgs field.

It is clear from Eq. 68 that the W fields will acquire a mass equal to $g\varphi_0/2$ from the term quadratic in the W fields, $(g^2/4)\varphi_0^2 W^\pm W_\mu^\pm$. The combination $g' B_\mu - g W_\mu^3$ will also have a mass. Thus, we “rotate” the B_μ and W_μ^3 fields to the fields Z_μ^0 for the weak neutral boson and A_μ for the photon so that the photon is massless.

$$\begin{pmatrix} Z_\mu^0 \\ A_\mu \end{pmatrix} = \begin{pmatrix} \sin \theta_W & -\cos \theta_W \\ \cos \theta_W & \sin \theta_W \end{pmatrix} \begin{pmatrix} B_\mu \\ W_\mu^3 \end{pmatrix}, \quad (69)$$

where

$$\cos \theta_W = g/\sqrt{g^2 + g'^2} \quad \text{and} \quad \sin \theta_W = g'/\sqrt{g^2 + g'^2}. \quad (70)$$

Upon substituting Eqs. 69 and 70 into Eq. 68, we find that the Z_μ^0 mass is $1/2 \varphi_0 \sqrt{g^2 + g'^2}$, so the ratio of the W and Z masses is

$$M_W/M_Z = \cos \theta_W. \quad (71)$$

Values for M_W and M_Z have recently been measured at the CERN proton-antiproton collider: $M_W = (80.8 \pm 2.7) \text{ GeV}/c^2$ and $M_Z = (92.9 \pm 1.6) \text{ GeV}/c^2$. The ratio M_W/M_Z calculated with these values agrees well with that given by Eq. 71. (The angle θ_W is usually expressed as $\sin^2 \theta_W$ and is measured in neutrino-scattering experiments to be $\sin^2 \theta_W = 0.224 \pm 0.015$.) The photon field A_μ does not appear in $\mathcal{L}_{\text{scalar}}$, so it does not become massive from spontaneous symmetry breaking. Note, also, that the $\pi^a(x)$ fields appear nowhere in the Lagrangian; they have been eaten by three weak vector bosons, which have become massive from the feast.

The next term in Eq. 56 is $\mathcal{L}_{\text{fermion}}$. Its form is analogous to Eqs. 39 and 40 for electrodynamics:

$$\mathcal{L}_{\text{fermion}} = i\bar{\Psi}\gamma^\mu D_\mu \Psi. \quad (72)$$

The physical problem is to assign the left- and right-handed fermions to multiplets of $SU(2)$; the assignments rely heavily on experimental data and are listed in “Particle Physics and the Standard Model.”

Our purpose here will be to write out Eq. 72 explicitly for the assignments.

Consider the electron and its neutrino. (The quark and remaining lepton contributions can be worked out in a similar fashion.) The left-handed components are assigned to a doublet and the right-handed components are singlets. (Since a neutral singlet has no weak charge, the right-handed component of the neutrino is invisible to weak, electromagnetic, or strong interactions. Thus, we can neglect it here, whether or not it actually exists.) We adopt the notation

$$\Psi_L = \begin{pmatrix} \nu_L \\ e_L^- \end{pmatrix} \quad \text{and} \quad \Psi_R = (e_R^-), \quad (73)$$

where L and R denote left- and right-handed. Then the explicit statement of Eq. 72 requires constructing D_μ for the left- and right-handed leptons.

$$\begin{aligned} \mathcal{L}_{\text{lepton}} = & i\bar{\Psi}_R \gamma^\mu (\partial_\mu + ig' B_\mu) \Psi_R \\ & + i\bar{\Psi}_L \gamma^\mu [\partial_\mu + \frac{i}{2} (g' B_\mu - g \tau_a W_\mu^a)] \Psi_L. \end{aligned} \quad (74)$$

The weak hypercharge of the right-handed electron is -2 so the coefficient of B_μ in the first term of Eq. 74 is $(-g'/2) \times (-2) = g'$. We leave it to the reader to check the rest of Eq. 74. The absence of a mass term is not an error. Mass terms are of the form $\bar{\psi}\psi = \bar{\psi}_L \psi_R + \bar{\psi}_R \psi_L$. Since Ψ_L is a doublet and $\bar{\psi}_R$ is a singlet, an electron mass term must violate the $SU(2) \times U(1)$ symmetry. We will see later that the electron mass will reappear as a result of modification of $\mathcal{L}_{\text{Yukawa}}$ due to spontaneous symmetry breaking.

The next task is exciting, because it will reveal how the vector bosons interact with the leptons. The calculation begins with Eq. 74 and requires the substitution of explicit matrices for $\tau_a W_\mu^a$, Ψ_R , and Ψ_L . We use the definitions in Eqs. 66, 69, and 73. The expressions become quite long, but the calculation is very straightforward. After simplifying some expressions, we find that $\mathcal{L}_{\text{lepton}}$ for the electron lepton and its neutrino is

$$\begin{aligned} \mathcal{L}_{\text{lepton}} = & i\bar{e}\gamma^\mu \partial_\mu e + i\bar{\nu}_L \gamma^\mu \partial_\mu \nu_L - e\bar{e}\gamma^\mu e A_\mu \\ & + \frac{g}{\sqrt{2}} (\bar{\nu}_L \gamma^\mu e_L W_\mu^+ + \bar{e}_L \gamma^\mu \nu_L W_\mu^-) \\ & - \frac{g^2}{2\sqrt{g^2 + g'^2}} [\tan^2 \theta_W (2\bar{e}_R \gamma^\mu e_R + \bar{e}_L \gamma^\mu e_L) - \bar{e}_L \gamma^\mu e_L] Z_\mu \\ & - \frac{1}{2} \sqrt{g^2 + g'^2} \bar{\nu}_L \gamma^\mu \nu_L Z_\mu. \end{aligned} \quad (75)$$

8 continued

The first two terms are the kinetic energies of the electron and the neutrino. (Note that $e = e_L + e_R$.) The third term is the electromagnetic interaction (cf. Eq. 40) with electrons of charge $-e$, where e is defined in Eq. 60. The coupling of A_μ to the electron current does not distinguish left from right, so electrodynamics does not violate parity. The fourth term is the interaction of the W^\pm bosons with the weak charged current of the neutrinos and electrons. Note that these bosons are blind to right-handed electrons. This is the reason for maximal parity violation in beta decay. The final terms predict how the weak neutral current of the electron and that of the neutrino couple to the neutral weak vector boson Z^0 .

If the left- and right-handed electron spinors are written out explicitly, with $e_L = \frac{1}{2}(1 - \gamma_5)e$, the interaction of the weak neutral current of the electron with the Z^0 is proportional to $\bar{e}\gamma^\mu[(1 - 4\sin^2\theta_W) - \gamma_5]eZ_\mu$. This prediction provided a crucial test of the standard model. Recall from Eq. 71 that $\sin^2\theta_W$ is very nearly $\frac{1}{4}$, so that the weak neutral current of the electron is very nearly a purely axial current, that is, a current of the form $\bar{e}\gamma^\mu\gamma_5e$. This crucial prediction was tested in deep inelastic scattering of polarized electrons and in atomic parity-violation experiments. The results of these experiments went a long way toward establishing the standard model. The tests also ruled out models quite similar to the standard model. We could discuss many more tests and predictions of the model based on the form of the weak currents, but this would greatly lengthen our discussion. The electroweak currents of the quarks will be described in the next section.

We now discuss the last term in Eq. 56, $\mathcal{L}_{\text{Yukawa}}$. In a locally symmetric theory with scalars, spinors, and vectors, the interactions between vectors and scalars, vector and spinors, and vectors and vectors are determined from the local invariance by replacing ∂_μ by D_μ . In contrast, $\mathcal{L}_{\text{Yukawa}}$, which is the interaction between the scalars and spinors, has the same form for both local and global symmetries:

$$\begin{aligned}\mathcal{L}_{\text{Yukawa}} &= G_Y \bar{\psi} \phi \psi \\ &= G_Y (\bar{\psi}_L \phi \psi_R + \bar{\psi}_R \phi^\dagger \psi_L).\end{aligned}\quad (76)$$

This form for $\mathcal{L}_{\text{Yukawa}}$ is rather schematic; to make it explicit we must

specify the multiplets and then arrange the component fields so that the form of $\mathcal{L}_{\text{Yukawa}}$ does not change under a local symmetry transformation.

Let us write Eq. 76 explicitly for the part of the standard model we have examined so far: ϕ is a complex doublet of scalar fields that has the form in the unitary gauge given by Eq. 65. The fermions include the electron and its neutrino. If the neutrino has no right-handed component, then it is not possible to insert it into Eq. 76. Since the neutrino has no mass term in $\mathcal{L}_{\text{lepton}}$, the neutrino remains massless in this theory. (If ν_R is included, then the neutrino mass is a free parameter.) The Yukawa terms for the electron are

$$\begin{aligned}\mathcal{L}_{\text{Yukawa}} &= G_Y \left[(\bar{\nu}_L, \bar{e}_L) \begin{pmatrix} 0 \\ (\rho + \phi_0)/\sqrt{2} \end{pmatrix} (e_R) \right. \\ &\quad \left. + (\bar{e}_R) (0, (\rho + \phi_0)/\sqrt{2}) \begin{pmatrix} \nu_L \\ e_L \end{pmatrix} \right] \\ &= \frac{1}{\sqrt{2}} G_Y \bar{e} e (\rho + \phi_0),\end{aligned}\quad (77)$$

where we have used the fact that $\bar{e}_L e_L = \bar{e}_R e_R = 0$, and $e = e_L + e_R$ is the electron Dirac spinor. Note that Eq. 77 includes an electron mass term,

$$m_e = \frac{1}{\sqrt{2}} G_Y \phi_0, \quad (78)$$

so the electron mass is proportional to the vacuum value of the scalar field. The Yukawa coupling is a free parameter, but we can use the measured electron mass to evaluate it. Recall that

$$M_W = \frac{g\phi_0}{2} = \frac{e\phi_0}{2 \sin \theta_W} \approx 81 \text{ GeV},$$

where $e^2/4\pi \approx 1/137$. This implies that $\phi_0 = 251 \text{ GeV}$. Since $m_e = 0.000511 \text{ GeV}$, $G_Y = 2.8 \times 10^{-6}$ for the electron. There are more than five Yukawa couplings, including those for the μ and τ leptons and the three quark doublets as well as terms that mix different quarks of the same electric charge. The standard model in no way determines the values of these Yukawa coupling constants. Thus, the study of fermion masses may turn out to have important hints on how to extend the standard model.



Quarks

Discovery of the fundamental fields of the strong interactions was not straightforward. It took some years to realize that the hadrons, such as the nucleons and mesons, are made up of subnuclear constituents, primarily quarks. Quarks originated from an effort to provide a simple physical picture of the "Eightfold Way," which is the SU(3) symmetry proposed by M. Gell-Mann and Y. Ne'eman to generalize strong isotopic spin. The hadrons could not be classified by the fundamental three-dimensional representations of this SU(3) but instead are assigned to eight- and ten-dimensional representations. These larger representations can be interpreted as products of the three-dimensional representations, which suggested to Gell-Mann and G. Zweig that hadrons are composed of constituents that are assigned to the three-dimensional representations: the *u* (up), *d* (down), and *s* (strange) quarks. At the time of their conception, it was not clear whether quarks were a physical reality or a mathematical trick for simplifying the analysis of the Eightfold-Way SU(3). The major breakthrough in the development of the present theory of strong interactions came with the realization that, in addition to electroweak and Eightfold-Way quantum numbers, quarks carry a new quantum number, referred to as color. This quantum number has yet to be observed experimentally.

We begin this lecture with a description of the Lagrangian of a strong-interaction theory of quarks formulated in terms of their color quantum numbers. Called quantum chromodynamics, or QCD, it is a Yang-Mills theory with local color-SU(3) symmetry in which each quark belongs to a three-dimensional color multiplet. The eight color-SU(3) generators commute with the electroweak SU(2) × U(1) generators, and they also commute with the generators of the Eightfold Way, which is a different SU(3). (Like SU(2), SU(3) is a recurring symmetry in physics, so its various roles need to be distinguished. Hence we need the label "color.") We conclude with a discussion of the weak interactions of the quarks.

The QCD Lagrangian. The interactions among the quarks are mediated by eight massless vector bosons (called gluons) that are required to make the SU(3) symmetry local. As we have already seen,

the assumption of local symmetry leads to a Lagrangian whose form is highly restricted. As far as we know, only the quark and gluon fields are necessary to describe the strong interactions, and so the most general Lagrangian is

$$\mathcal{L}_{\text{QCD}} = -\frac{1}{4} F_{\mu\nu}^a F_{\mu\nu}^a + i \sum_i \bar{\psi}_i \gamma^\mu D_\mu \psi_i + \sum_{i,j} \bar{\psi}_i M_{ij} \psi_j, \quad (79)$$

where

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g_s f_{abc} A_\mu^b A_\nu^c. \quad (80)$$

The sum on *a* in the first term is over the eight gluon fields A_μ^a . The second term represents the coupling of each gluon field to an SU(3) current of the quark fields, called a color current. This term is summed over the index *i*, which labels each quark type and is independent of color. Since each quark field ψ_i is a three-dimensional column vector in color space, D_μ is defined by

$$D_\mu \psi_i = \partial_\mu \psi_i - \frac{1}{2} i g_s A_\mu^a \lambda_a \psi_i, \quad (81)$$

where λ_a is a generalization of the three 2-by-2 Pauli matrices of SU(2) to the eight 3-by-3 Gell-Mann matrices of SU(3), and g_s is the QCD coupling. Thus, the color current of each quark has the form $\bar{\psi} \lambda_a \gamma^\mu \psi$. The left-handed quark fields couple to the gluons with exactly the same strength as the right-handed quark fields, so parity is conserved in the strong interactions.

The gluons are massless because the QCD Lagrangian has no spinless fields and therefore no obvious possibility of spontaneous symmetry breaking. Of course, if motivated for experimental reasons, one can add scalars to the QCD Lagrangian and spontaneously break SU(3) to a smaller group. This modification has been used, for example, to explain the reported observation of fractionally charged particles. The experimental situation, however, still remains murky, so it is not (yet) necessary to spontaneously break SU(3) to a smaller group. For the remainder of the discussion, we assume that QCD is not spontaneously broken.

The third term in Eq. 79 is a mass term. In contrast to the electroweak theory, this mass term is now allowed, even in the absence of spontaneous symmetry breaking, because the left- and right-handed quarks are assigned to the same multiplet of SU(3). The numerical coefficients M_{ij} are the elements of the quark mass matrix; they can connect quarks of equal electric charge. The \mathcal{L}_{QCD} of Eq. 79 permits us to redefine the QCD quark fields so that $M_{ij} = m_i \delta_{ij}$. The

9 continued

mass matrix is then diagonal and each quark has a definite mass, which is an eigenvalue of the mass matrix. We will reappraise this situation below when we describe the weak currents of the quarks.

After successfully extracting detailed predictions of the electroweak theory from its complicated-looking Lagrangian, we might be expected to perform a similar feat for the \mathcal{L}_{QCD} of Eq. 79 without too much difficulty. This is not possible. Analysis of the electroweak theory was so simple because the couplings g and g' are always small, regardless of the energy scale at which they are measured, so that a classical analysis is a good first approximation to the theory. The quantum corrections to the results in Note 8 are, for most processes, only a few percent.

In QCD processes that probe the *short-distance* structure of hadrons, the quarks inside the hadrons interact weakly, and here the classical analysis is again a good first approximation because the coupling g_s is small. However, for Yang-Mills theories in general, the renormalization group equations of quantum field theory require that g_s increases as the momentum transfer decreases until the momentum transfer equals the masses of the vector bosons. Lacking spontaneous symmetry breaking to give the gluons mass, QCD contains no mechanism to stop the growth of g_s , and the quantum effects become more and more dominant at larger and larger distances. Thus, analysis of the *long-distance* behavior of QCD, which includes deriving the hadron spectrum, requires solving the full quantum theory implied by Eq. 79. This analysis is proving to be very difficult.

Even without the solution of \mathcal{L}_{QCD} , we can, however, draw some conclusions. The quark fields ψ_i in Eq. 79 must be determined by experiment. The Eightfold Way has already provided three of the quarks, and phenomenological analyses determine their masses (as they appear in the QCD Lagrangian). The mass of the u quark is nearly zero (a few MeV/c^2), the d quark is a few MeV/c^2 heavier than the u , and the mass of the s quark is around $300 \text{ MeV}/c^2$. If these results are substituted into Eq. 79, we can derive a beautiful result from the QCD Lagrangian. In the limit that the quark mass differences can be ignored, Eq. 79 has a global $\text{SU}(3)$ symmetry that is identical to the Eightfold-Way $\text{SU}(3)$ symmetry. Moreover, in the limit that the u , d , and s masses can be ignored, the left-handed u , d ,

and s quarks can be transformed by one $\text{SU}(3)$ and the right-handed u , d , and s quarks by an independent $\text{SU}(3)$. Then QCD has the "chiral" $\text{SU}(3) \times \text{SU}(3)$ symmetry that is the basis of current algebra. The sums of the corresponding $\text{SU}(3)$ generators of chiral $\text{SU}(3) \times \text{SU}(3)$ generate the Eightfold-Way $\text{SU}(3)$. Thus, the QCD Lagrangian incorporates in a very simple manner the symmetry results of hadronic physics of the 1960s. The more recently discovered c (charmed), b (bottom), and t (top) quarks are easily added to the QCD Lagrangian. Their masses are so large and so different from one another that the $\text{SU}(3)$ and $\text{SU}(3) \times \text{SU}(3)$ symmetries of the Eightfold-Way and current algebra cannot be extended to larger symmetries. (The predictions of, say, $\text{SU}(4)$ and chiral $\text{SU}(4) \times \text{SU}(4)$ do not agree well with experiment.)

It is important to note that the quark masses are undetermined parameters in the QCD Lagrangian and therefore must be derived from some more complete theory or indicated phenomenologically. The Yukawa couplings in the electroweak Lagrangian are also free parameters. Thus, we are forced to conclude that the standard model alone provides no constraints on the quark masses, so they must be obtained from experimental data.

The mass term in the QCD Lagrangian (Eq. 79) has led to new insights about the neutron-proton mass difference. Recall that the quark content of a neutron is udd and that of a proton is uud . If the u and d quarks had the same mass, then we would expect the proton to be more massive than the neutron because of the electromagnetic energy stored in the uu system. (Many researchers have confirmed this result.) Since the masses of the u and d quarks are arbitrary in both the QCD and the electroweak Lagrangians, they can be adjusted phenomenologically to account for the fact that the neutron mass is $1.293 \text{ MeV}/c^2$ greater than the proton mass. This experimental constraint is satisfied if the mass of the d quark is about $3 \text{ MeV}/c^2$ greater than that of the u quark. In a way, this is unfortunate, because we must conclude that the famous puzzle of the n - p mass difference will not be solved until the standard model is extended enough to provide a theory of the quark masses.

Weak Currents. We turn now to a discussion of the weak currents of the quarks, which are determined in the same way as the weak currents of the leptons in Note 8. Let us begin with just the u and d quarks. Their electroweak assignments are as follows: the left-handed components u_L and d_L form an $\text{SU}(2)$ doublet with $Y = 1/3$, and the right-handed components u_R and d_R are $\text{SU}(2)$ singlets with $Y = 4/3$

and $-2/3$, respectively (recall Eq. 55).

The steps followed in going from Eq. 73 to Eq. 75 will yield the electroweak Lagrangian of quarks. The contribution to the Lagrangian due to interaction of the weak neutral current $J_\mu^{(nc)}$ of the u and d quarks with Z^0 is

$$\mathcal{L}^{(nc)} = \frac{e}{\sin \theta_W \cos \theta_W} J_\mu^{(nc)} Z_\mu^0, \quad (82)$$

where

$$J_\mu^{(nc)} = \left(\frac{1}{2} - \frac{2}{3} \sin^2 \theta_W \right) \bar{u}_L \gamma_\mu u_L - \frac{2}{3} \sin^2 \theta_W \bar{u}_R \gamma_\mu u_R \\ + \left(-\frac{1}{2} + \frac{1}{3} \sin^2 \theta_W \right) \bar{d}_L \gamma_\mu d_L + \frac{1}{3} \sin^2 \theta_W \bar{d}_R \gamma_\mu d_R. \quad (83)$$

The reader will enjoy deriving this result and also deriving the contribution of the weak charged current of the quarks to the electroweak Lagrangian. Equation 83 will be modified slightly when we include the other quarks.

So far we have emphasized in Notes 8 and 9 the construction of the QCD and electroweak Lagrangians for just one lepton-quark "family" consisting of the electron and its neutrino together with the u and d quarks. Two other lepton-quark families are established experimentally: the muon and its neutrino along with the c and s quarks and the τ lepton and its neutrino along with the t and b quarks. Just like $(\nu_e)_L$ and e_L , $(\nu_\mu)_L$ and μ_L and $(\nu_\tau)_L$ and τ_L form weak-SU(2) doublets; e_R , μ_R and τ_R are each SU(2) singlets with a weak hypercharge of -2 . Similarly, the weak quantum numbers of c and s and of t and b echo those of u and d : c_L and s_L form a weak-SU(2) doublet as do t_R and b_L . Like u_R and d_R , the right-handed quarks c_R , s_R , t_R , and b_R are all weak-SU(2) singlets.

This triplification of families cannot be explained by the standard model, although it may eventually turn out to be a critical fact in the development of theories of the standard model. The quantum numbers of the quarks and leptons are summarized in Tables 2 and 3 in "Particle Physics and the Standard Model."

All these quark and lepton fields must be included in a Lagrangian that incorporates both the electroweak and QCD Lagrangians. It is quite obvious how to do this: the standard model Lagrangian is

simply the sum of the QCD and electroweak Lagrangians, except that the terms occurring in both Lagrangians (the quark kinetic energy terms $i\bar{\psi}_i \gamma^\mu \partial_\mu \psi_i$ and the quark mass terms $\bar{\psi}_i M_{ij} \psi_j$) are included just once. Only the mass term requires comment.

The quark mass terms appear in the electroweak Lagrangian in the form $\mathcal{L}_{\text{Yukawa}}$ (Eq. 77). In the electroweak theory quarks acquire masses only because SU(2) \times U(1) is spontaneously broken. However, when there are three quarks of the same electric charge (such as d , s , and b), the general form of the mass terms is the same as in Eq. 79, $\bar{\psi}_i M_{ij} \psi_j$, because there can be Yukawa couplings between d and s , d and b , and s and b . The problem should already be clear: when we speak of quarks, we think of fields that have a definite mass, that is, fields for which M_{ij} is diagonal. Nevertheless, there is no reason for the fields obtained directly from the electroweak symmetry breaking to be mass eigenstates.

The final part of the analysis takes some care: the problem is to find the most general relation between the mass eigenstates and the fields occurring in the weak currents. We give the answer for the case of two families of quarks. Let us denote the quark fields in the weak currents with primes and the mass eigenstates without primes. There is freedom in the Lagrangian to set $u = u'$ and $c = c'$. If we do so, then the most general relationship among d , s , d' , and s' is

$$\begin{pmatrix} d' \\ s' \end{pmatrix} = \begin{pmatrix} \cos \theta_C & -\sin \theta_C \\ \sin \theta_C & \cos \theta_C \end{pmatrix} \begin{pmatrix} d \\ s \end{pmatrix}. \quad (84)$$

The parameter θ_C , the Cabibbo angle, is not determined by the electroweak theory (it is related to ratios of various Yukawa couplings) and is found experimentally to be about 13° . (When the b and t ($=t'$) quarks are included, the matrix in Eq. 84 becomes a 3-by-3 matrix involving four parameters that are evaluated experimentally.) The correct weak currents are then given by Eq. 83 if all quark families are included and primes are placed on all the quark fields. The weak currents can be written in terms of the quark mass eigenstates by substituting Eq. 84 (or its three-family generalization) into the primed version of Eq. 83. The ratio of amplitudes for $s \rightarrow u$ and $d \rightarrow u$ is $\tan \theta_C$; the small ratio of the strangeness-changing to non-strangeness-changing charged-current amplitudes is due to the smallness of the Cabibbo angle. It is worth emphasizing again that the standard model alone provides no understanding of the value of this angle. \odot